arXiv:1703.03474v1 [physics.class-ph] 8 Mar 2017

Numerical analysis of the location of spectral maxima of polarization components of synchrotron radiation in the classical theory

A. S. Loginov [1], A. D. Saprykin [2]

March 13, 2017

Abstract

Shift of higher harmonic peak in the spectrum of synchrotron radiation with increasing energy of the radiating particle is investigated by numerical method. Calculations were carried out for the $\nu^{(max)} \leqslant 100$ for all polarization components.

1 Introduction

At present, the theory of synchrotron radiation is a fairly well developed area of theoretical physics. Its main elements are described in monographs (e.g. [1–8]) and numerous articles

Spectral distribution of SR has one physically important feature, namely increasing energy of radiating particle leads to shift of spectral maximum to higher harmonics. This feature was predicted in works [9, 10] and has been confirmed by numerous experiments.

It's convenient to represent energy E of radiating particle by relativistic factor $\gamma\,(\gamma \geqslant 1)$, or quantity $\beta\,(0 \leqslant \beta < 1)$

$$\gamma = \frac{E}{m_0 c^2} = \frac{1}{\sqrt{1-\beta^2}}\,, \quad \beta = \frac{v}{c}\,, \tag{1. 1}$$

where m_0 – the charge rest mass, v is the charge orbital motion speed, c – the speed of lite. With respect to notation of rotation frequency of radiating particle by ω_0, spectrum of SR has frequencies $\omega = \omega_0 \nu$, where whole number $\nu = 1,\ 2,\ 3, \ldots$ – number of radiated harmonic.

For nonrelativistic particle which corresponds to the condition $\beta \ll 1$ (equivalently $\gamma \sim 1$) maximum of SR spectrum falls on first harmonics $\nu^{(max)} = 1$. With increasing energy of the particle spectral maximum shifts sequentially to 2,3,4... harmonics and for ultra-relativistic particle (corresponding to $\gamma \gg 1$) asymptotic estimate $\nu^{(max)} \sim \gamma^3$ holds (It was shown in [9, 10]). This estimate is correct for polarization component of SR.

Exact values of β (or γ) which lead to shift of maximum in SR spectral distribution from harmonic ν to harmonic ν+1 can be carried out by numerical calculations. within the framework of classical theory such calculations have been carried out in the work [11] for $\nu \leqslant 15$ for the total radiated power and the linear polarization component of SR. This paper presents the results of numerical analysis for $\nu \leqslant 100$ and all polarization component of SR.

[1]Tomsk State University, Russia. E-mail: as_loginov@phys.tsu.ru

[2]Tomsk State University, Russia. E-mail: alexx.saprykin.phys@gmail.com

2 Spectral distribution of SR polarization component in upper half-space

To put a numerical analysis of the problem considered in this article we present here some well-known expression of the physical characteristics SR in the classical theory which can be found in [1–8].

The spectral - angular distribution of the radiation power of the SR polarization component can be represented by

$$W_s = W\sum_{\nu=1}^{\infty}\int_0^{\pi} f_s(\beta;\,\nu,\,\theta)\sin\theta d\theta. \qquad (2.\ 1)$$

Here, the following notation is used: θ is the angle between the control magnetic field strength and the radiation field pulse, W – the total radiated power of unpolarized radiation, which can be written as

$$W = \frac{2}{3}\frac{ce^2}{R^2}(\gamma^2-1)^2 = \frac{2e^4H^2(\gamma^2-1)}{3m_0^2c^3}\,, \qquad (2.\ 2)$$

where e – the particle charge;; R – orbit radius;; H – is the control field strength. The index s numbers the polarization components:: $s=2$ corresponds to σ - component of linear polarization; $s=3$ corresponds to π - component of linear polarization; $s=1$ corresponds to right-hand circular polarization; $s=-1$ – corresponds to left-hand circular polarization;; $s=0$ corresponds to the power of unpolarized radiation. The functions $f_s(\beta;\,\nu,\,\theta)$ have the form

$$f_2(\beta;\,\nu,\,\theta) = \frac{3\nu^2}{2\gamma^4}J_\nu'^{\,2}(x);\quad f_3(\beta;\,\nu,\,\theta) = \frac{3\nu^2}{2\gamma^4}\frac{\cos^2\theta}{\beta^2\sin^2\theta}J_\nu^2(x);$$

$$f_{\pm1}(\beta;\,\nu,\,\theta) = \frac{3\nu^2}{4\gamma^4}\left[J_\nu'(x)\pm\varepsilon\frac{\cos\theta}{\beta\sin\theta}J_\nu(x)\right]^2;\quad x=\nu\beta\sin\theta;\quad \varepsilon=-\frac{e}{|e|};$$

$$f_0(\beta;\,\nu,\,\theta) = f_2(\beta;\,\nu,\,\theta)+f_3(\beta;\,\nu,\,\theta) = f_1(\beta;\,\nu,\,\theta)+f_{-1}(\beta;\,\nu,\,\theta)\,. \qquad (2.\ 3)$$

Here, $J_\nu(x)$ and $J_\nu'(x)$ are the Bessel functions and their derivatives. In what follows, the case of an electron is considered, which corresponds to $\varepsilon=1$.

It is well known [1–7] that the angle range $0\leqslant\theta<\pi/2$ (this range will be called the upper half-space) is dominated by right-hand circular polarization, and the angle range $\pi/2<\theta\leqslant\pi$ (this range will be called the lower half-space) is dominated by left-hand circular polarization (exact quantitative characteristics of SR properties were first obtained in [12–15]). However, if we integrate in (2. 1)) over θ $(0\leqslant\theta\leqslant\pi)$ then the differences in the spectral distribution of right- and left-hand circular polarizations disappear. To reveal these differences, a new form of expression (2. 1)) has been proposed in [16] It can be represented as

$$W_s = W_s^{(+)}+W_s^{(-)}\,,\quad W_s^{(\pm)} = W\sum_{\nu=1}^{\infty}F_s^{(\pm)}(\beta;\,\nu)\,;$$

$$F_s^{(+)}(\beta;\,\nu) = \int_0^{\pi/2} f_s(\beta;\,\nu,\,\theta)\sin\theta d\theta\,,\quad F_s^{(-)}(\beta;\,\nu) = \int_{\pi/2}^{\pi} f_s(\beta;\,\nu,\,\theta)\sin\theta d\theta\,, \qquad (2.\ 4)$$

and it suffices to study the properties of functions $F_s^{(+)}(\beta;\,\nu)$ due to the evident relations

$$F_s^{(-)}(\beta;\,\nu) = F_s^{(+)}(\beta;\,\nu)\,,\quad s=0,\,2,\,3;$$

$$F_1^{(-)}(\beta;\,\nu) = F_{-1}^{(+)}(\beta;\,\nu)\,,\quad F_{-1}^{(-)}(\beta;\,\nu) = F_1^{(+)}(\beta;\,\nu)\,. \qquad (2.\ 5)$$

Integration over θ in the upper half-space $0 \leqslant \theta \leqslant \pi/2$ in (2. 4) can be carried out exactly, yielding the expressions

$$F_2^{(+)}(\beta;\,\nu) = \frac{3\nu}{4\,\gamma^{\,4}\beta^3}\left[2\beta^2 J'_{2\nu}(2\nu\beta) + \beta^2\int_0^{2\nu\beta} J_{2\nu}(x)dx - 2\nu\beta\int_0^{2\nu\beta}\frac{J_{2\nu}(x)}{x}dx\right],$$

$$F_3^{(+)}(\beta;\,\nu) = \frac{3\nu}{4\,\gamma^{\,4}\beta^3}\left[2\nu\beta\int_0^{2\nu\beta}\frac{J_{2\nu}(x)}{x}dx - \int_0^{2\nu\beta} J_{2\nu}(x)dx\right],$$

$$F_0^{(+)}(\beta;\,\nu) = F_2^{(+)}(\beta;\,\nu) + F_3^{(+)}(\beta;\,\nu) = \frac{3\nu}{4\,\gamma^{\,4}\beta^3}\left[2\beta^2 J'_{2\nu}(2\nu\beta) - (1-\beta^2)\int_0^{2\nu\beta} J_{2\nu}(x)dx\right],$$

$$F_{\pm1}^{(+)}(\beta;\,\nu) = \frac{1}{2}F_0^{(+)}(\beta;\,\nu) \pm \frac{3\nu J_\nu^{\,2}(\nu\beta)}{4\,\gamma^{\,4}\beta^2}\,. \qquad (2.\ 6)$$

3 Numerical analysis of the shift of the maximum in the SR spectrum with increasing the energy of the radiating particle

Expressions (ref B6) make possible to translate into formal language the task of analyzing the shift of SR spectral maximum at change of radiating particle energy E.

As is known, in the non-relativistic limit (which corresponds to $\beta \to 0$ and $\gamma \to 1$) maxima of any components $W_s^{(+)}$ in the spectral distribution occur in first harmonic ($\nu_s^{(max)} = 1$). With increasing the energy of particle (increase of β or, respectively, γ) maximum in the spectrum of $W_s^{(+)}$ goes to the second harmonic.

Obviously, the value of $\beta = \beta_1^{(s)}$, which is carried out at the maximum shift in the spectrum of $W_s^{(+)}$ to the second harmonic is the root of the equation

$$F_s^{(+)}(\beta;\,1) = F_s^{(+)}(\beta;\,2)\,. \qquad (3.\ 1)$$

So for $0 \leqslant \beta < \beta_1^{(s)}$ (or $1 \leqslant \gamma < \gamma_1^{(s)}$) the maximum in spectrum of $W_s^{(+)}$ falls on the first harmonic

With increasing β (or γ) maximum in spectrum of $W_s^{(+)}$ shift to second harmonic. The value $\beta = \beta_2^{(s)}$, corresponding to this shift, is root of equation

$$F_s^{(+)}(\beta;\,2) = F_s^{(+)}(\beta;\,3)\,. \qquad (3.\ 2)$$

So for $\beta_1^{(s)} \leqslant \beta < \beta_2^{(s)}$ (or $\gamma_1^{(s)} \leqslant \gamma < \gamma_2^{(s)}$) maximum in spectrum of $W_s^{(+)}$ falls on second harmonic.

In general, for a fixed s determining the sequence of numbers $\beta = \beta_\nu^{(s)}$ as the roots of equations

$$F_s^{(+)}(\beta;\,\nu) = F_s^{(+)}(\beta;\,\nu+1)\,, \qquad (3.\ 3)$$

It can be found that in energy range

$$\beta_{\nu-1}^{(s)} \leqslant \beta < \beta_\nu^{(s)} \quad \left(\text{respectively } \gamma_{\nu-1}^{(s)} \leqslant \gamma < \gamma_\nu^{(s)}\right) \qquad (3.\ 4)$$

maximum in spectrum of $W_s^{(+)}$ falls on harmonic ν.

The following Table for $\nu \leqslant 100$ for each polarization s $\beta_\nu^{(s)}$ and $\gamma_\nu^{(s)}$ are given. It should be noted that following inequalities are correct

$$\gamma_\nu^{(3)} > \gamma_\nu^{(1)} > \gamma_\nu^{(0)} > \gamma_\nu^{(2)} > \gamma_\nu^{(-1)} \quad \left(\text{respectively } \beta_\nu^{(3)} > \beta_\nu^{(1)} > \beta_\nu^{(0)} > \beta_\nu^{(2)} > \beta_\nu^{(-1)}\right). \qquad (3.\ 5)$$

It is also easy to see from Table, that since $\nu_{-1}^{(max)}(\beta) = 8$ inequalities, obtained in [16] for ultra-relativistic particle, hold.

In the ultra-relativistic case ($1 \ll \gamma$) for $\nu_s^{(max)} = 1$ asymptotic formulas are known.

$$\nu_s^{(max)} \approx a_s \gamma^3\,, \qquad (3.\ 6)$$

where a_s – numbers, approximate values of ones are equal (see [16]) to

$$a_3 \approx 0,215881\,,\quad a_1 \approx 0,372875\,,\quad a_0 \approx 0,428718\,,\quad a_2 \approx 0,503287\,,\quad a_{-1} \approx 0,783608\,. \quad (3.\ 7)$$

If $\nu_s^{(max)} = 100$ value corresponding γ determined by the asymptotic formulas (3. 6), they may differ from the table of values in no more than fourth place.

Table 1: Boundary values of energy (corresponding to $\beta_\nu^{(s)}$) for spectral maximum shift

ν	$\beta_\nu^{(2)}$	$\gamma_\nu^{(2)}$	$\beta_\nu^{(3)}$	$\gamma_\nu^{(3)}$	$\beta_\nu^{(0)}$	$\gamma_\nu^{(0)}$	$\beta_\nu^{(-1)}$	$\gamma_\nu^{(-1)}$	$\beta_\nu^{(1)}$	$\gamma_\nu^{(1)}$
1	0,71493	1,43020	0,88148	2,11768	0,76354	1,54856	0,58762	1,23588	0,78799	1,62420
2	0,80568	1,68819	0,91747	2,51376	0,83681	1,82649	0,72085	1,44281	0,85442	1,92461
3	0,84815	1,88768	0,93305	2,77982	0,87113	2,03643	0,78352	1,60936	0,88498	2,14761
4	0,87340	2,05334	0,94249	2,99182	0,89174	2,20971	0,82060	1,74978	0,90325	2,33037
5	0,89036	2,19655	0,94902	3,17254	0,90572	2,35920	0,84538	1,87209	0,91563	2,48749
6	0,90265	2,32359	0,95390	3,33211	0,91594	2,49175	0,86325	1,98108	0,92469	2,62652
7	0,91203	2,43835	0,95773	3,47611	0,92378	2,61147	0,87682	2,07979	0,93164	2,75195
8	0,91946	2,54340	0,96082	3,60802	0,93002	2,72110	0,88753	2,17033	0,93719	2,86671
9	0,92552	2,64056	0,96340	3,73018	0,93513	2,82252	0,89621	2,25418	0,94173	2,97282
10	0,93056	2,73115	0,96557	3,84428	0,93941	2,91712	0,90343	2,33243	0,94553	3,07177
11	0,93483	2,81619	0,96745	3,95154	0,94304	3,00594	0,90953	2,40593	0,94876	3,16465
12	0,93851	2,89644	0,96908	4,05295	0,94618	3,08979	0,91477	2,47533	0,95156	3,25232
13	0,94171	2,97253	0,97052	4,14924	0,94892	3,16932	0,91932	2,54115	0,95400	3,33546
14	0,94453	3,04495	0,97180	4,24104	0,95133	3,24504	0,92331	2,60383	0,95616	3,41461
15	0,94704	3,11411	0,97295	4,32884	0,95348	3,31738	0,92686	2,66370	0,95808	3,49022
16	0,94928	3,18037	0,97399	4,41306	0,95541	3,38669	0,93002	2,72106	0,95980	3,56266
17	0,95130	3,24400	0,97493	4,49404	0,95715	3,45327	0,93287	2,77615	0,96136	3,63224
18	0,95313	3,30525	0,97579	4,57209	0,95873	3,51737	0,93545	2,82919	0,96277	3,69924
19	0,95480	3,36433	0,97658	4,64745	0,96018	3,57922	0,93780	2,88035	0,96406	3,76388
20	0,95633	3,42142	0,97730	4,72036	0,96150	3,63899	0,93995	2,92980	0,96525	3,82636
21	0,95774	3,47668	0,97797	4,79099	0,96272	3,69687	0,94192	2,97766	0,96634	3,88684
22	0,95904	3,53025	0,97860	4,85954	0,96385	3,75298	0,94374	3,02407	0,96735	3,94549
23	0,96025	3,58226	0,97918	4,92614	0,96489	3,80746	0,94543	3,06911	0,96829	4,00244
24	0,96137	3,63281	0,97972	4,99092	0,96587	3,86043	0,94700	3,11290	0,96916	4,05780
25	0,96241	3,68200	0,98023	5,05401	0,96678	3,91198	0,94846	3,15550	0,96997	4,11168
26	0,96339	3,72992	0,98071	5,11552	0,96763	3,96220	0,94982	3,19701	0,97074	4,16417
27	0,96431	3,77664	0,98116	5,17554	0,96843	4,01118	0,95110	3,23748	0,97145	4,21537
28	0,96517	3,82225	0,98158	5,23416	0,96918	4,05899	0,95230	3,27699	0,97213	4,26535

Table 1: Boundary values of energy (corresponding to $\beta_\nu^{(s)}$) for spectral maximum shift

ν	$\beta_\nu^{(2)}$	$\gamma_\nu^{(2)}$	$\beta_\nu^{(3)}$	$\gamma_\nu^{(3)}$	$\beta_\nu^{(0)}$	$\gamma_\nu^{(0)}$	$\beta_\nu^{(-1)}$	$\gamma_\nu^{(-1)}$	$\beta_\nu^{(1)}$	$\gamma_\nu^{(1)}$
29	0,96598	3,86680	0,98198	5,29146	0,96989	4,10570	0,95343	3,31557	0,97277	4,31417
30	0,96675	3,91035	0,98236	5,34750	0,97055	4,15137	0,95450	3,35330	0,97337	4,36191
31	0,96747	3,95296	0,98272	5,40237	0,97119	4,19606	0,95551	3,39020	0,97394	4,40862
32	0,96816	3,99467	0,98306	5,45611	0,97179	4,23981	0,95646	3,42633	0,97447	4,45436
33	0,96881	4,03554	0,98339	5,50879	0,97236	4,28268	0,95737	3,46172	0,97499	4,49918
34	0,96943	4,07560	0,98370	5,56045	0,97290	4,32470	0,95823	3,49642	0,97547	4,54311
35	0,97002	4,11489	0,98399	5,61115	0,97342	4,36593	0,95905	3,53045	0,97594	4,58620
36	0,97058	4,15345	0,98427	5,66093	0,97391	4,40639	0,95983	3,56384	0,97638	4,62850
37	0,97112	4,19131	0,98454	5,70982	0,97438	4,44611	0,96057	3,59663	0,97681	4,67004
38	0,97163	4,22850	0,98480	5,75787	0,97483	4,48514	0,96128	3,62884	0,97721	4,71084
39	0,97213	4,26505	0,98505	5,80511	0,97526	4,52350	0,96196	3,66049	0,97760	4,75095
40	0,97260	4,30099	0,98529	5,85158	0,97567	4,56123	0,96261	3,69161	0,97797	4,79039
41	0,97305	4,33634	0,98552	5,89731	0,97607	4,59833	0,96324	3,72222	0,97833	4,82919
42	0,97348	4,37113	0,98574	5,94232	0,97645	4,63485	0,96384	3,75234	0,97867	4,86737
43	0,97390	4,40537	0,98595	5,98665	0,97681	4,67080	0,96441	3,78200	0,97900	4,90496
44	0,97430	4,43909	0,98616	6,03032	0,97716	4,70620	0,96496	3,81120	0,97931	4,94198
45	0,97468	4,47231	0,98635	6,07335	0,97750	4,74108	0,96550	3,83996	0,97962	4,97845
46	0,97505	4,50505	0,98654	6,11576	0,97783	4,77546	0,96601	3,86830	0,97991	5,01439
47	0,97541	4,53731	0,98673	6,15759	0,97814	4,80934	0,96650	3,89624	0,98020	5,04982
48	0,97576	4,56913	0,98690	6,19884	0,97845	4,84276	0,96698	3,92379	0,98047	5,08476
49	0,97609	4,60051	0,98707	6,23954	0,97874	4,87571	0,96744	3,95096	0,98074	5,11923
50	0,97641	4,63148	0,98724	6,27970	0,97903	4,90823	0,96788	3,97776	0,98099	5,15323
51	0,97672	4,66203	0,98740	6,31935	0,97930	4,94033	0,96831	4,00422	0,98124	5,18679
52	0,97703	4,69219	0,98756	6,35850	0,97957	4,97201	0,96873	4,03033	0,98148	5,21992
53	0,97732	4,72197	0,98771	6,39716	0,97982	5,00329	0,96913	4,05611	0,98171	5,25264
54	0,97760	4,75138	0,98785	6,43535	0,98007	5,03418	0,96952	4,08157	0,98194	5,28495
55	0,97788	4,78043	0,98800	6,47309	0,98031	5,06470	0,96990	4,10672	0,98215	5,31686
56	0,97814	4,80914	0,98813	6,51038	0,98055	5,09486	0,97027	4,13157	0,98237	5,34840
57	0,97840	4,83751	0,98827	6,54724	0,98078	5,12466	0,97062	4,15612	0,98257	5,37957
58	0,97865	4,86555	0,98840	6,58369	0,98100	5,15413	0,97097	4,18039	0,98277	5,41038
59	0,97890	4,89327	0,98852	6,61972	0,98121	5,18325	0,97130	4,20439	0,98297	5,44085
60	0,97913	4,92068	0,98865	6,65537	0,98142	5,21206	0,97163	4,22811	0,98315	5,47097
61	0,97936	4,94779	0,98877	6,69063	0,98163	5,24055	0,97195	4,25158	0,98334	5,50077
62	0,97959	4,97461	0,98888	6,72551	0,98182	5,26873	0,97225	4,27479	0,98352	5,53024
63	0,97981	5,00114	0,98900	6,76004	0,98202	5,29661	0,97255	4,29776	0,98369	5,55941
64	0,98002	5,02740	0,98911	6,79420	0,98220	5,32420	0,97285	4,32048	0,98386	5,58827
65	0,98023	5,05338	0,98922	6,82802	0,98239	5,35151	0,97313	4,34297	0,98402	5,61683
66	0,98043	5,07910	0,98932	6,86151	0,98256	5,37855	0,97341	4,36523	0,98419	5,64511
67	0,98062	5,10457	0,98943	6,89466	0,98274	5,40531	0,97368	4,38727	0,98434	5,67310
68	0,98082	5,12978	0,98953	6,92750	0,98291	5,43181	0,97394	4,40909	0,98450	5,70082
69	0,98100	5,15475	0,98963	6,96002	0,98307	5,45806	0,97420	4,43070	0,98464	5,72827
70	0,98119	5,17948	0,98972	6,99223	0,98323	5,48405	0,97445	4,45210	0,98479	5,75546
71	0,98136	5,20398	0,98981	7,02415	0,98339	5,50980	0,97469	4,47330	0,98493	5,78240
72	0,98154	5,22825	0,98991	7,05577	0,98355	5,53531	0,97493	4,49430	0,98507	5,80908

Table 1: Boundary values of energy (corresponding to $\beta_\nu^{(s)}$) for spectral maximum shift

ν	$\beta_\nu^{(2)}$	$\gamma_\nu^{(2)}$	$\beta_\nu^{(3)}$	$\gamma_\nu^{(3)}$	$\beta_\nu^{(0)}$	$\gamma_\nu^{(0)}$	$\beta_\nu^{(-1)}$	$\gamma_\nu^{(-1)}$	$\beta_\nu^{(1)}$	$\gamma_\nu^{(1)}$
73	0,98171	5,25229	0,99000	7,08711	0,98370	5,56059	0,97517	4,51511	0,98521	5,83553
74	0,98187	5,27612	0,99008	7,11817	0,98384	5,58564	0,97539	4,53573	0,98534	5,86173
75	0,98204	5,29974	0,99017	7,14896	0,98399	5,61047	0,97562	4,55616	0,98547	5,88770
76	0,98220	5,32315	0,99025	7,17948	0,98413	5,63508	0,97583	4,57642	0,98560	5,91345
77	0,98235	5,34635	0,99033	7,20974	0,98427	5,65948	0,97605	4,59650	0,98572	5,93897
78	0,98250	5,36936	0,99042	7,23975	0,98440	5,68367	0,97626	4,61640	0,98584	5,96427
79	0,98265	5,39217	0,99049	7,26950	0,98453	5,70765	0,97646	4,63614	0,98596	5,98936
80	0,98280	5,41479	0,99057	7,29901	0,98466	5,73144	0,97666	4,65571	0,98608	6,01424
81	0,98294	5,43722	0,99065	7,32828	0,98479	5,75502	0,97686	4,67512	0,98619	6,03892
82	0,98308	5,45947	0,99072	7,35732	0,98491	5,77842	0,97705	4,69437	0,98631	6,06339
83	0,98322	5,48154	0,99079	7,38612	0,98503	5,80163	0,97724	4,71347	0,98642	6,08767
84	0,98335	5,50344	0,99086	7,41470	0,98515	5,82465	0,97742	4,73241	0,98652	6,11176
85	0,98349	5,52516	0,99093	7,44305	0,98527	5,84750	0,97760	4,75121	0,98663	6,13565
86	0,98361	5,54672	0,99100	7,47119	0,98538	5,87016	0,97778	4,76986	0,98673	6,15937
87	0,98374	5,56810	0,99107	7,49912	0,98550	5,89265	0,97795	4,78836	0,98683	6,18290
88	0,98387	5,58933	0,99114	7,52683	0,98561	5,91497	0,97812	4,80672	0,98693	6,20625
89	0,98399	5,61039	0,99120	7,55435	0,98571	5,93713	0,97829	4,82495	0,98703	6,22943
90	0,98411	5,63130	0,99126	7,58165	0,98582	5,95912	0,97845	4,84303	0,98713	6,25243
91	0,98422	5,65206	0,99133	7,60877	0,98592	5,98095	0,97861	4,86099	0,98722	6,27527
92	0,98434	5,67266	0,99139	7,63568	0,98603	6,00262	0,97877	4,87882	0,98731	6,29794
93	0,98445	5,69312	0,99145	7,66241	0,98613	6,02413	0,97892	4,89651	0,98740	6,32045
94	0,98456	5,71343	0,99151	7,68895	0,98622	6,04550	0,97908	4,91408	0,98758	6,36500
95	0,98467	5,73360	0,99157	7,71530	0,98632	6,06671	0,97923	4,93153	0,98758	6,36500
96	0,98478	5,75363	0,99162	7,74148	0,98642	6,08777	0,97937	4,94885	0,98775	6,40892
97	0,98489	5,77351	0,99168	7,76747	0,98651	6,10869	0,97952	4,96606	0,98775	6,40892
98	0,98499	5,79327	0,99173	7,79329	0,98660	6,12947	0,97966	4,98314	0,98784	6,43066
99	0,98509	5,81289	0,99179	7,81894	0,98669	6,15011	0,97980	5,00012	0,98792	6,45225
100	0,98519	5,83238	0,99184	7,84442	0,98678	6,17061	0,97993	5,01697	0,98800	6,47370

Acknowledgements

The work of Loginov and Saprykin is supported by Tomsk State University Competitiveness Improvement Program.

References

[1] A. A. Sokolov, I. M. Ternov. Synchrotron Radiation. Akademie - Verlag, 1968, 202 p.

[2] Herman Winick, S. Doniach. Synchrotron Radiation Research. Plenum Press, 1980, 754 p.

[3] A. A. Sokolov, I. M. Ternov, C. W. Kilmister. Radiation from Relativistic Electrons. American Institute of Physics, 1986, 312 p.

[4] V. G. Bagrov et al. Synchrotron Radiation Theory and Its Development. (World Scientific, Singapore, New Jersey, London, 1999), 447 p.

[5] P. J. Duke. Synchrotron Radiation: Production and Properties. Oxford University Press, 2000, 251 p.

[6] V. G. Bagrov, G. S. Bisnovaty – Kogan, V. A. Bordovitsyn, A. V. Borisov, O. F. Dorofeev, V. Ya. Épp, Yu. L. Pivovarov, O. V. Shorokhov, V. Ch. Zhukovsky. Radiation Theory of Relativistic Particles. Ed. V. A. Bordovitsyn. Moscow, Fizmatlit. 2002, 576 p. (In Russian).

[7] Helmut Wiedemann. Synchrotron Rdiation. Springer, 2003, 274 p.

[8] Albert Hofmann. The Physics of Synchrotron Radiation. Cambridge University Press, 2004, 323 p.

[9] L. Arzimovich, I. Pomeranchuk. The Radiation of Fast Electrons in the Magnetic Field. Journal of Physics. 1945, v. IX, No 4, p. 267-276.

[10] L. Arzimovich, I. Pomeranchuk. The Radiation of Fast Electrons in the Magnetic Field. Soviet Physics JETP. 1946, v. 16, No 5, p. 379-389. (In Russian).

[11] V. G. Bagrov, G. F. Kopytov, G. K. Razina, V. B. Tlyachev. Numerical Analysis of Spectral Distribution of Synchrotron Radiation (Classical Theory). Dep. VINITI 12.11.85, No 7927 - B 85, 13 p.(In Russian).

[12] . V. G. Bagrov, M. V. Dolzhin, V. B. Tlyachev, A. T. Jarovoi. Evolution of the Angular Distribution of Circularly Polarized Synchrotron Radiation with Charge Energy. Russian Physics Journal. 2004, v. 47, No. 4, pp. 414-423, doi:10.1023/B:RUPJ.0000042770.97166.87

[13] V. G. Bagrov, D. M. Gitman, V. B. Tlyachev, A. T. Jarovoi. Evolution of Angular Distribution of Polarization Components for Synchrotron Radiation under Changes of Particle Energy. International Summer School: Recent Problems in Field Theory. V. 4. Petrov School. Kazan, June - July 2003. Ed. A. V. Aminovain, Kazan. Heter, 2004, pp. 9-24. (In Russian).

[14] V. G. Bagrov, D. M. Gitman, V. B. Tlyachev, A. T. Jarovoi. Evolution of Angular Distribution of Polarization Components for Synchrotron Radiation under Changes of Particle Energy. International Conference: Particle Physics in Laboratory, Space and Universe. Moscow State University, Faculty Physics, August 2003. Ed. A. I. Studenikin, World Scientific. New Jersey, London, Singapor, 2005, pp.355-362.

[15] V. G. Bagrov, D. M. Gitman, V. B. Tlyachev, A. T. Jarovoi. New Theoretical Results in Synchrotron Radiation. Nuclear Instruments & Methods in Physics Research B. 2005, v. 240, No. 3, pp. 638-645, doi:10.1016/j.nimb.2005.03.286

[16] V. G. Bagrov, D. M. Gitman,A. D. Levin, A. S. Loginov, A. D. Saprykin. Effective Spectrum Width of the Synchrotron Radiation. The European Physical Journal. C. 2015, v. 75, No. 11, 555, 6 p., doi:10.1140/epjc/s10052-015-3798-6